

## STORAGE AND DISPOSAL

**STORAGE:** Store in original container and place in a locked storage area.

**CONTAINER DISPOSAL:** Do not reuse container. Wrap and put in trash.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**NOTE TO USER:** This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER:** May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Do not breathe spray mist. Skin sensitizer, wash thoroughly after handling.

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED**—Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED**—Remove victim to fresh air. Apply artificial respiration if indicated.

**IF ON SKIN**—Remove contaminated clothing. Wash affected areas with soap and water.

**IF IN EYES**—Flush eyes with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of liability incurred shall be the purchase price paid by the user or buyer.

Product freezing temperature 32°F. Satisfactory utility is restored after thawing and mixing of product. Repeated 0-10° freezing-thawing may impair utility.

\*Chlorothalonil U.S. Patent 3,290,353 and No. 3,331,735.

# Hi-Yield®

## Daconil® LAWN, VEGETABLE AND FLOWER FUNGICIDE



- Unique Fungicide for Lawn, Vegetable, Flowers & Ornamentals
- Controls Brown Patch, Dollar Spot, Leaf Spot and Certain other Fungi

#### ACTIVE INGREDIENT:

Chlorothalonil	
(tetrachloroisoph thalonitrile)*	12.5%
INERT INGREDIENTS:	87.5%
TOTAL	100.0%

**KEEP OUT OF REACH  
OF CHILDREN**

## DANGER

**SEE SIDE PANEL FOR  
ADDITIONAL CAUTIONS**

**NET CONTENTS ONE QUART (32 FL. OZS.)**

Hi-Yield® Chemical Company  
Bonham, Texas 75418  
EPA Est. No. 7401-TX-1  
EPA Reg. No. 7401-65-34911  
23M-23-21H



#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN  
A MANNER INCONSISTENT WITH ITS LABELING.

We recommend this highly effective new fungicidal formulation, because it will make your lawn and garden disease control program easier.

#### SHAKE WELL BEFORE USE:

**LAWN**—For the control of Brown Patch, Dollar Spot, Leaf Spot (Blight, Going-out, Fading-out and Melting-out), Gray Leaf Spot, Stem Rust on Bluegrass and Copper Spot: Mix 1 pint in 5-15 gallons of water and apply to 500 sq. ft. of lawn. The amount of water should be sufficient to provide complete coverage. DO NOT mow or water grass after treatment until the grass is thoroughly dry and preferably not for 24 hours after treatment. Start spray application in Spring before diseases appear and repeat at 7-14 day intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Do not graze treated areas. Do not feed clippings to livestock.

**VEGETABLES AND MELONS**—Unless otherwise specified, mix 3 tablespoons in 5-15 gallons of water and apply to 500 sq. ft. of garden area. Repeat at 7 to 10 day intervals or as necessary to maintain control. Carefully follow application schedule under individual crops.

**BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER**—For Alternaria Leaf Spot (S.E. States only) and Downy Mildew. Apply after transplant, emergence of seeded crop or when conditions favor disease development.

**CARROTS**—For Early and Late Blight. Begin when disease threatens.

**POTATOES**—For Botrytis Vine Rot, Early and Late Blight. Begin when plants are 6 inches high and disease threatens.

**PUMPKIN, SQUASH (Summer and Winter), WATERMELON**—For Anthracnose and Downy Mildew. Use 6 tablespoons per 5-15 gallons of water per 500 sq. ft. for Cercospora Leaf Spot, Gum Stem, Leaf Blight and Scab. Begin treatment in the first leaf stage.

**TOMATOES**—For Anthracnose, Early and Late Blights, Gray Leaf Spot, Gray Leaf Mold. Use 4 tablespoons in 5-15 gallons of water per 500 sq. ft. Begin when disease threatens.

**MELONS**—(Cantaloupe, Honeydew, Muskmelon)—For Anthracnose, Downy Mildew. Use 6 tablespoons in 5-15 gallons of water per 500 sq. ft. For Powdery Mildew (except in S.W. States), Cercospora Leaf Spot, Gummy Stem Blight, Leaf Blight, Scab. Begin when plants are in first true leaf stage or conditions favor disease development.

**CHRYSANTHEMUMS**—For control of Botrytis Blight (Leaf Spot).

**IRIS**—For the control of Botrytis Blight (Blossom Blight) and Iris Leaf Spot.

**GERANIUMS**—For the control of Botrytis Blight (Leaf Spot).

**ZINNIA**—For the control of Powdery Mildew: Mix 9 tablespoons in 1 gallon of water. Spray plants on both sides of foliage at the drip point. Penetrate dense foliage. Apply at first sign of disease and repeat at 7-14 day intervals throughout growing season. Make application at 7 day schedule during period of wet weather or severe disease conditions.

**ROSES**—For control of Black Spot, Blossom Blight: Mix 6 tablespoons in one gallon of water.