

Havahart
Caring Control for Pets and Wildlife

Deer & rabbit Deer Away[®]

- ♥ Repels deer for up to 3 months.
- ♥ Creates an odor barrier to protect garden vegetables.*
- ♥ Protects landscape plantings, ornamental trees and shrubs.
- ♥ Concentrate: Makes 5 Gallons.

ACTIVE INGREDIENT: BY WT.
Purified Whole Egg Solids..... 23.15%
OTHER INGREDIENTS: 76.85%
TOTAL: 100.00%



FOR RESIDENTIAL USE ONLY
DO NOT SPRAY DIRECTLY ON PLANTS INTENDED FOR HUMAN CONSUMPTION.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements and First Aid.

NET CONTENTS: 1 Gallon (3.79 l)

www.havahart.com



Havahart
Caring Control for Pets and Wildlife

deer & rabbit Deer Away[®] CONCENTRATE

General Information:
Before applying, read and understand all label instructions. SHAKE WELL BEFORE MIXING! Do not spray directly onto plants intended for human consumption. This product can be applied using all types of hand or power sprayers. Read and carefully follow the instructions for the type of sprayer you use. Promptly clean all equipment after each use. Do not mix or apply this product with equipment that has been used for herbicidal applications. Properly dilute and spray this product on treatment area. Do not apply excessively to seedlings.

Precautionary Statements:
Hazards to Humans and Domestic Animals:
CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

First Aid:
If in eyes:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.
If on skin or clothing:
- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact 1-800-858-7378 for emergency medical treatment advice.

Environmental Hazards:
For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinseate.

Directions For Use:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions:
For home & garden use only. **IMPORTANT:** This product contains egg solids, which may cause allergic reactions in some individuals. Not for direct use on plants intended for human consumption (i.e. food plants). Do not allow spray to drift onto plants intended for human consumption. Use this product to repel black-tailed deer (mule deer), white-tailed deer and Roosevelt elk that browse on uproot conifer seedlings. This product can be used to repel white-tailed deer from view. Scotch pine, arborvitae, Douglas fir, flowering crabapple, hennock, honeysuckle,

stelia, rhododendron, landscape gardens and plantings, ornamental shrubs, garden vegetables, and fruit and citrus trees. Use this product to repel rabbits from ornamentals, landscape plantings and garden vegetables. Before applying, read specific application directions and use cautions below. Store at room temperature. Protect from heat and freezing. Freezing makes this product unusable. This product will stain structural surfaces. Do not apply to or allow spray to drift onto siding, roofing, decks, fences, driveways, painted surfaces or other structural material.

Mixing and Dilution Instructions:
Note: Container is not filled to capacity but is filled by weight.
Use before using: This product must be mixed with water to produce ready-to-use spray. Before mixing, read and understand label instructions.
Use promptly after mixing: Mixed solution will spoil quickly above 40°F. Do not store mixed solution longer than one day. Carefully determine the quantity required for immediate application.

1. Before diluting, thoroughly shake or stir contents of concentrate container to completely resuspend all solids.
2. In a suitable mixing container, mix 1 part concentrate with 4 parts warm water.
3. Pour dilute spray into sprayer and use promptly. Stir or shake the spray immediately before use to prevent settling in bottom of sprayer. Run agitator on large tank sprayers. If sprayer is to be reused, clean it with cold water immediately after use.
Seal tightly: If saving part of unmixed concentrate for use later, close container tightly to prevent evaporation.

Application Directions:
Apply only in diluted form as specified. Do not apply excessively to seedlings.

Weather Conditions: Do not apply at temperatures below 40°F, or in wet, windy conditions. Apply only to dry (non-food) plants, and only under conditions favoring prompt drying of spray (lightly breezy, mild dry weather with low relative humidity). If spray is used on wet (non-food) plants in damp, cold foggy or misty weather or if precipitation or freezing occurs before spray can dry protection may be partially or completely lost.

Care of Equipment: Promptly discard leftover spray according to label directions. Rinse spraying equipment thoroughly with cold water immediately after use.

New Conifer Seedlings: New seedlings to be protected from winter browsing must be dormant and frost hardy before treatment. Once planted, retreat seedlings every 6 weeks during root establishment or as needed if previous treatment appears to be fading. Note: Browning and drop of old needles may occur on some evergreen species after using this product.

Established Conifer Seedlings: Treat each established seedling to be protected. Use a pressurized hand-held or backpack garden sprayer. Hold nozzle of spray wand about 10" vertically above tip of terminal shoot. Spray the terminal and first whorl to run-off. Treat each seedling to be protected.



Yew, Scotch Pine, Arborvitae, Douglas Fir, Ornamental Crabapple, Hennock, Honeysuckle, Azalea and Rhododendron.

To protect against winter browsing, apply repellent upon onset of browsing. If browsing extends beyond two months, seedlings may be re-treated. Apply only to vegetation that is within animals' reach. Apply to the point of run-off. Reapply as needed if the previous treatment appears to be fading.

To protect against spring browsing of new growth, apply repellent upon onset of browsing, but preferably after bud break and before new shoots exceed 1" in length. Do not re-treat new growth if spring browsing reaches unacceptable proportions before bud break. Repellent may be applied to unopened buds, but must then be re-applied after new growth averages 1" in length if browsing still persists. Apply to terminal to point of run-off. For large trees, apply this product only to lateral branches that are within animals' reach. Reapply as needed if the previous treatment appears to be fading. Note: Browning and drop of old needles may occur on some evergreen species after using this product.

Ornamental Shrubs: Do not apply during bud break or flower bloom, as foliage injury and petal drop may occur. Apply only to areas of vegetation that are within animals' reach. Apply to the point of run-off. Reapply as needed if the previous treatment appears to be fading.

Non-Bearing Fruit and Citrus Trees: For use on non-bearing fruit & citrus trees only. Use when trees are dormant, not when actively growing. Do not apply during bud break or flower bloom, as foliage injury and petal drop may occur. Apply only to vegetation that is within animals' reach. Apply to the point of run-off. Reapply as needed if the previous treatment appears to be fading.

Landscape Gardens: Do not apply to landscape plantings, during bud break or flower bloom, as foliage injury or petal drop may occur. Apply to areas of vegetation that are within animals' reach. Apply to the point of run-off. Reapply as needed if the previous treatment appears to be fading.

Vegetable Gardens: This product is not for direct application to plants intended for human consumption (i.e. food plants). For use as a perimeter spray around the area where the food plants are grown. To create a repellent barrier, apply this product to soil, ground cover, bark mulch, perimeter vegetation and non-food vegetation adjacent to food plants. Do not allow spray to drift from application site onto any part of the plant that is intended for human consumption. Reapply as needed if the previous treatment appears to be fading. *Not registered for use as a perimeter spray in California.

Storage and Disposal:
Do not contaminate water, food or feed by storage and disposal.
Pesticide Storage: Store only in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or in an approved waste disposal facility.
Container Disposal:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.
If Partly Filled: Contact your local solid waste agency or 1-800-CLERK for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured by:
Woodstream

Woodstream Corporation, 69 North Locust Street, Lititz, PA 17543
For information, call (800) 800-1819 or visit www.havahart.com

EPA Reg. No. 50932-15
EPA Est. No. C3416-MN-001

Made in USA Model # 5212

0905 © Copyright 2004 Woodstream Corporation