



Specimen Label

- May be used on Livestock Animals and Livestock Premises
- Kills Bed Bugs
- Kills Bees and Wasps [Yellow Jackets, Hornets, and Paper Wasps]

ACTIVE INGREDIENTS:

Pyrethrins, a botanical insecticide.....	1.00%
OTHER INGREDIENTS	99.00%
	100.00%

0.01 lbs. active ingredient per lbs. of Evergreen® Pyrethrum Dust



KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE

USE RESTRICTIONS AND PRECAUTIONS

- Apply this product only as specified on the label.
- Aerial applications are prohibited.
- Applications with power duster equipment are prohibited.
- Do not apply this product in a way that will contact workers or other persons, either directly or indirectly through drift.
- Only protected handlers may enter treated area during application.
- Do not allow adults, children or pets to enter the treated area until dusts have settled.
- For use outdoors, do not apply directly to or near water, storm drains or drainage ditches. Do not apply when windy. To prevent product run-off, do not over-water the treated area or apply when heavy rain is expected.
- Do not apply this product in a way that will contact adults, children or pets, either directly or indirectly through drift.
- Remove or cover exposed food and drinking water before application.
- In Homes remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use. Do not use in serving areas while food is exposed.

- In Food Areas of Food Handling Establishments: Avoid contamination of food or feedstuffs. During applications, cover or remove food. Cover food processing surfaces and equipment or clean after treatment with a suitable detergent and rinse with potable water before use. Do not use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation.
- In Schools, do not apply to classrooms when in use.
- Do not use in aircraft.
- Remove pets, birds and cover fish aquariums before application.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until dusts have settled.

Kills Exposed Insects and Related Pests including:

Adelgids	Brown Spruce	Crickets	Foreign Grain Beetles	Larger Grain Borers	Pinworms	Sowbugs
Alfalfa Weevil	Longhorn Beetle	Cross-striped	Forest Tent Caterpillar	Leafhoppers	Plant Bugs	Spider Beetles
Almond Moth	Budworms	Cabbageworm	Fruit Flies	Leafminers	Plum Curculio	Spider Mites
American Black Flour Beetle	Bugs	Cucumber Beetles	Fruittree Leafroller	Leafrollers	Ponderosa Pine Needle Miner	Spruce Mite (adult)
Angoumois Grain Moths	Cadelles	Cutworms (exposed)	Fungus Beetles	Leaftiers	Poultry Lice	Squarenecked Grain Beetles
Ants	California Oakworm	Dark Mealworms	Fungus Gnats	Lesser Grain Beetles	Powder Post Beetles	Stable Flies
Aphids	Cankerworms	Darkling Beetles (lesser meal worm)	German Cockroaches	Lesser Grain Borers	Psocids	Stalk Borers
Apple Maggot	Carpet Beetles	Deer Tick	Giant Root Aphid	Lesser Meal Worm	Psyllids	Stink Bugs
Armyworms	Carriion Beetles	Depressed Flour Beetles	Glassy Winged Sharpshooter	Lesser Peach Tree Borer	Red Flour Beetles	Stored Product Pests
Asian Cerambycid Beetles	Caterpillars	Dermestid Beetles	Grain Mites	Longheaded Flour Beetles	Red Gum Lerp Psyllid	Subtropical Pine Tip Moth
Asian Lady Beetle	Catorama Beetles	Diamondback Larvae	Granary Insects	Loopers	Red Horned Grain Beetles	Sycamore Leaf Beetle
Asian Pine Tip Moths	Centipedes	Dingy Cutworm Adult	Granary Weevils	Lygus Bugs	Red Lily Beetles	Tarnished Plant Bug
Bagworm	Cereal Beetles	Douglas Fir Midge	Grape Leaf Skeletonizer	Maize Weevils	Red Oak Borer	Tent Caterpillar
Barn Flies	Cheese Mites	Douglas Fir Tussock Moth	Grape Leafhopper	Mealy Bugs	Redbanded Leafroller	Thrips
Bat Bugs	Cheese Skippers	Drain Flies	Grasshoppers	Mediterranean Flour Moths	Rice Flour Beetles	Tobacco Moths
Bean Beetles	Chicken Mites	Dried Fruit Beetles	Green Fruit Worm	Merchant Grain Beetles	Rice Moths	Trogoderma Beetles
Bean Weevils	Chocolate Moths	Drugstore Beetles	Green Peach Aphid	Mexican Bean Beetle	Rice Weevils	Two-Banded Beetles
Bed Bugs	Chrysanthemum Leaf Miners	Earwigs	Greenbugs	Mexican Grain Beetles	Root Weevils	Vegetable Leaf Miners
Bees	Cicadas	Eastern Tent Caterpillar	Greenhouse Thrips	Millers	Rose Chafer	Vinegar Flies
Beet Armyworms	Cigarette Beetle	Elmleaf Beetle	Ground Beetles	Millipedes	Royal Palm Bugs	Warehouse Beetles
Beetles	Clearwing Borer	Emerald Ash Borer	Gypsy Moth (adults & larvae)	Mimosa Webworm	Rusty Grain Beetles	Wasps [Yellow Jackets, Hornets, and Paper Wasps]
Birch Leafminer	Cloves Moths	European Grain Moths	Hairy Fungus Beetles	Mole Crickets	Saw Flies	Waterbugs
Black Carpet Beetles	Clover Mites	European Pine Shoot Moth	Hairy Spider Beetles	Moth Flies	Saw-Toothed Grain Beetle	Webworms
Black Fungus Beetles	Clover Weevil	European Pine Tip Moth	Harlequin Bug	Murmidius Beetles	Scale (Crawlers)	Weevils
Black Turpentine Beetles	Cluster Flies	Face Flies	Heliiothis spp.	Mushroom Flies	Seedbugs	Western Pine Tip Moth
Black Vine Weevil	Cocoa Bean Moths	Fall Cankerworms	Hessian Fly (adult)	Nantucket Pine Tip Moth	Silverfish	White Apple Leafroller
Blacklegged Tick Nymphs	Coffee Bean Weevils	Fall Webworm	Hide Beetles	Northern Fowl Mites	Skipper Flies	White Marked Spider Beetles
Blister Beetles	Cone Beetles	Fannia Flies	Hornworm	Onion Maggot	Slenderhorned Flour Beetles	Whiteflies
Blow Flies	Cone Worms	Field Crickets	Inchworm	Palmetto Bug	Small Flying Moths	Wireworms
Booklice	Confused Flour Beetle	Firebrats	Indian Meal Moth	Peach Twig Borer	Small Fruit Flies	Yellow Mealworms
Bottle Flies	Conifer Spider Mite (adult)	Fireworms	Iris Borer	Pear Psylla	Smalleyed Flour Beetles	
Boxelder Bugs	Corn Sap Beetles	Flat Grain Beetles	Japanese Beetle	Pillbugs	Southern Pine Beetles	
Broadhorned Flour Beetles	Cottonwood Borer	Flea Beetles	Katydids	Pine Bark Beetles	Southern Red Mite (adult)	
Broadnosed Grain Weevils	Cottonwood Leaf Beetles	Fleas	Khapra Beetles	Pine Engraver		
Brown Spider Beetles	Cranbury Girdlers	Flies	Lace Bugs	Pink Scavenger Caterpillars		
	Crane Flies	Flying Moths	Large Black Flour Beetles			
			Larger Cabinet Beetles			

READ ENTIRE LABEL BEFORE EACH USE

For most effective results, follow directions for specific use areas.

Avoid contamination of food and feedstuffs.

Application Methods:

Apply Evergreen® Pyrethrum Dust using a dust applicator, a shaker can, dust gun, puffer or other bulbous duster, hand pump duster, hand operated rotary duster, static charge duster, or other appropriate hand operated duster.

INDOOR USE SITES:

ONLY APPLY AS A SPOT TREATMENT TO AREAS NO GREATER THAN 2 FEET BY 2 FEET PER ROOM.

INDOOR AREAS: For use in and around indoor areas:

Animal Areas	Coffee Houses	Egg Processing Plants	Goat Houses	Livestock Holding Areas	Poultry Processing Plants	Tobacco Factories
Animal Quarters	Coffee Shops	Egg Storage Facilities	Grain Elevators	Loafing Sheds	Rabbit Houses	Tobacco Warehouses
Apartments	Commercial Buildings	Factories	Grain Handling Equipment	Meat Markets	Rabbit Processing Plants	Tool Sheds
Asian Markets	Condominiums	False Ceilings	Grain Harvesting Equipment	Meat Packaging Plants	Railroad Cars	Trains
Attics	Convenience Stores	Farm Structures	Grain Mills	Meat Packing Plants	Research Animal Areas	Transportation Equipment (except Aircraft)
Bakeries	Crawl Spaces	Feed Mills	Grain Storage	Mess Halls	Restaurants	Tree Nuts Storage Areas
Barns	Dairies	Flour Mills	Granaries	Milk Houses	Rice Mills	Truck Trailers
Basements	Dairy and Livestock Barns	Food Handling Establishments	Greenhouses	Milk Processing Facilities	Schools	Vegetable Packing Facilities
Beef Barns	Dairy Food Processing	Food Preparation Facilities	Grocery Stores	Milk Rooms	Seed Storage	Vegetable Processing Facilities
Beverage Plants	Day Care Centers	Food Processing Facilities	Hog Barns	Milk Sheds	Shade Houses	Vegetable Storage Facilities
Bins	Delicatessens	Food Quarantine Areas	Homes	Milking Rooms	Sheep Holding Areas	Vegetable Storage Facilities
Bottling Plants	Department Stores	Food Service Facilities	Horse Stables	Mills	Ships	Veterinary Clinics
Brandy Storage Warehouses	Diners	Food Storage Facilities	Horse Stables and Barns	Motels	Ships	Veterinary Hospitals
Breweries	Distilleries	Freight Containers	Horse Trailers	Mushroom Caves	Spice Markets	Voids in Attics
Brewery Warehouses	Dog Houses	Fruit Packing Sheds	Hospitals	Mushroom Processing Plants	Stable Bins	Wall Voids
Buildings Housing Animals	Dog Sleeping Quarters	Fruit Processing Plants	Hotels	Office Buildings	Stables	Warehouses
Buildings Housing Poultry	Drains & Sewers	Fruit Quarantine Areas	Houses	Other Public Buildings	Storage Areas	Wheat Mills
Cabinets	Dried Flower Warehouses	Fruit Storage Warehouses	Indoor Eating Facilities	Packaged Food Storage	Storage Bins (areas)	Wholesale Stores
Cafeterias	Dried Food Products in Processing	Garbage Bins	Industrial Installations	Pantries	Storages	Wine Cellars
Calving Barns	Dried Food Products in Storage	Garbage Cans	Industrial Plants	Peanut Processing Plants	Stored Product Areas	Wine Storage Warehouses
Canneries	Dried Fruit Processing Plants	Garbage Compactors	Institution Dining Areas	Peanut Warehouses	Stored Products	Wine Tasting Rooms
Cat Sleeping Quarters	Dried Fruit Warehouses	Garbage Trucks	Institutions	Pet Bedding	Stored Seed Warehouses	Wineries
Cattle Barns	Dried Fruit Warehouses	Goat Holding Areas	Kennels	Pet Carriers	Stores	Zoos
Certified Organic Facilities	Dumpsters		Liveries	Pet Food Plants	Supermarkets	
Cheese Caves	Dwellings		Livestock Barns	Pet Sleeping Areas	Swine Houses	
Citrus Packing Houses			Livestock Hauling Equipment	Poultry Houses	Taverns	
					Tea Houses	
					Textile Mills	

APPLICATIONS WITHIN FOOD/FEED AREAS

Food and/or Feed areas include areas for receiving, storage (dry, cold, frozen or raw), packaging (canning, bottling, wrapping, boxing and grinding), preparing, edible waste storage and enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups). Serving areas are places where prepared foods are served. Do not apply when food or feed processing facility is in operation or foods are exposed. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment.

APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS ARE NOT PERMITTED, OTHER THAN AS A CRACK AND CREVICE TREATMENT.

Using a duster, apply at the rate of 1.5 ounces of Evergreen® Pyrethrum Dust per 50 square feet directly into cracks and crevices, in expansion joints between different elements of construction, under or between equipment bases and the floor, beneath appliances, into wall voids, motor housing, junction boxes or switch boxes, conduits, behind and under cabinets or drawers, in and around drains, around drain pipes, plumbing and utility entrances, hollow equipment legs and other sites where listed insects can enter or hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Non-food/feed areas of food/feed handling establishments are garbage rooms, lavatories, floor drains (to sewers, entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Stored Product Pests in Non-Food/Non-Feed Areas:

To aid in the control of stored product pests in listed areas. Apply Evergreen® Pyrethrum Dust at the rate of 16 to 24 ounces per 1,000 square feet [4 to 6 ounces per 250 square feet] of surface area. Clean up waste materials, infested materials, dust, dirt, and any other debris. Thoroughly treat floors, walls, shelves, drawers, cabinets and other surfaces in bins, storage and handling areas, around cartons or containers and other areas where these insects tend to congregate. Apply 1 ounce per 50 square feet of surface area into cracks and crevices and other difficult to reach areas where insects may hide. Apply in voids or cracks and crevices in pantries and other infested areas within the structure. When treating for Indian Meal Moth, which tend to alight on vertical surfaces, apply a thin visible layer of Evergreen® Pyrethrum Dust to walls and other vertical surfaces in affected area.

For Use in Storage Sites:

This product can be used in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment and other listed sites as a clean up prior to using them for storage. All of the infested stored commodity accumulations and debris should be removed and disposed, with special attention to material lodged in the cracks and crevices. After sanitation measures have been employed, dust all areas prior to use for storage. Apply Evergreen® Pyrethrum Dust, at the rate of 16 to 24 ounces per 1,000 square feet [4 to 6 ounces per 250 square feet] of surface area.

Apply dust to floors, ceilings and partition boards of bins, paying particular attention to cracks and crevices.

For packaged food warehouses, treat potential points of insect entry, as well as harborage sites in, on and around wooden and plastic pallets, pallet-racking systems, loading docks, scales, around cartons.

INDOOR TREATMENTS

ONLY APPLY AS A SPOT TREATMENT TO AREAS NO GREATER THEN 2 FEET BY 2 FEET PER ROOM.

Crawling Insects:

Apply Evergreen® Pyrethrum Dust, at the rate of 1.5 ounce per 50 square feet of surface area, into voids, cracks and crevices or other places where insects are traveling or harboring. Apply a continuous visible film underneath and behind stairs, behind and beneath sinks, stoves, refrigerators and other appliances, around garbage cans, cabinets. Treat on and around and behind moldings, shelving, baseboards, doorsills, doorframes, floors, around drains, pipes and pipe openings, plumbing, bookcases, storage, utility installations and other areas where insects hide or travel. For severe infestations apply behind cabinets and walls, in crawl spaces and attics at the rate of 16 - 20 ounces per 1,000 square feet [4 to 5 ounces per 250 square feet].

Fleas, Ticks, and Mites:

For Cat Fleas apply Evergreen® Pyrethrum Dust only as a direct treatment. Treat on and around animal bedding, sleeping quarters, kennel, furniture, floor and floor coverings and all areas where these pests inhabit. Treat cracks and crevices behind around window and door frames, wall cracks moldings, shelving, baseboards, pipe openings and all other cracks. Treat the entire inside surface of doghouses. Do not treat dogs and/or cats with this product. For Ticks, apply Evergreen® Pyrethrum Dust, at the rate of 1 ounce per 70 square feet in places where the insects are traveling or harboring.

Bed Bugs:

Take bed apart. Apply Evergreen® Pyrethrum Dust at a rate of 2 oz. per 50 square feet of surface area into joints, channels, cracks and crevices on bed frame, headboards and footboards. Wall mounted headboards and their hardware should be removed and treated. If hollow, square or round tubing, see that the interior framework is well dusted. Mattresses and box springs may be dusted, especially tufts, folds and edges. All furniture and furniture accessories in the room must be treated. Pull carpeting from tack strips and treat under the baseboard, sill plate and under the carpet tack strips. Treat under, behind, and into cracks and crevices of decorative moldings. Apply a continuous visible film to other potential harborage sites: table and wall mounted lamps, wall crevices, electrical boxes, outlet plate covers, switch plate covers, mirrors, picture frames and backs, door jams, moldings and hinges, window ledges, sills, drapery rods, curtains, blinds and mounting hardware, closet shelving and hardware, picture frame moldings and other crack and crevices in the room should be treated. If Bed Bugs are present, treat with an Insecticide approved for use on Bed Bugs.

Ants:

To aid in the control of listed ants apply Evergreen® Pyrethrum Dust at the rate of 1 oz. per 50 square feet. Treat door frames, around window frames, behind, on and around moldings, cabinets, pipe openings, utility openings, shelves, ant trails, hills, wall voids, cracks and crevices and other points of entry into premises. Combine interior treatments with exterior perimeter application of Evergreen® Pyrethrum Dust.

Powder Post Beetles*:

To aid in protecting against insect entry. Apply to distribute Evergreen® Pyrethrum Dust through insect entry points, crawl holes into attics and crawl spaces of buildings, at a rate of 16 - 20 ounces per 1,000 square feet [4 to 5 ounces per 250 square feet]. All vents and blowers on the windward side should be closed so that all target surfaces can be treated. Contact all wood surfaces. In case of localized infestations, probe to locate galleries and drill holes to inject Evergreen® Pyrethrum Dust into the galleries and void areas. If a heavy infestation is suspected, a conventional, industry accepted techniques and treatments should be employed.

*The purpose of such applications of Evergreen® Pyrethrum Dust is to kill workers or winged reproductive forms that may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not for use against termites. For active termite infestations, use conventional, industry accepted techniques and treatments.

INDOOR AND OUTDOOR TREATMENTS

For Bees and Wasps [Yellow Jackets, Hornets, and Paper Wasps]:

It is advisable to treat bee and wasp [yellow jacket, hornet, and paper wasp] nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Use a hand duster with extension tubes if necessary. Thoroughly dust the nest, entrance and surrounding areas. For best results, check nests carefully one or two days after treatment to ensure complete kill. Remove and destroy nest to prevent emergence of newly hatched insects. If removal is not possible, retreat the nest if necessary. For nests in wall voids, drill a hole in the area of the nest, apply dust, and reseal.

Asian Lady Beetles and Boxelder Bugs:

To control or repel these insects, as well as Clover Mites, Cluster Flies and Elmleaf Beetles, inject Evergreen® Pyrethrum Dust at the rate of 1 oz. per 50 square feet into cracks and crevices and/or voids where these insects may be entering the premise, harboring or hibernating. Apply in crawl spaces and attics at the rate of 16 - 24 ounces per 1,000 square feet [4 to 6 ounces per 250 square feet]. On building exterior, treat underneath wood fascia, behind siding, cracks around windows, doors, siding, wall voids, utility pipes and entry points, behind chimneys, around foundations and foundation cracks, patios and deck joints. When treating for Asian Lady Beetles apply Evergreen® Pyrethrum Dust in, on and around areas with abrupt color contrasts on buildings (shutters, drainpipes, etc), southwest-facing or sun-exposed sides of buildings, and vertical corners of buildings.

For Use on Landscape Plants, Indoors and Outdoors:

Ornamental Trees, Shrubs, Flowers and Foliage Plants: Including: African Violet, Ageratum, Aster, Azalea, Begonia, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Cineraria, Chrysanthemum, Coleum, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Fern, Ficus, Foliage Plants, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Holly, Hyacinth, Hydrangea, Iris, Lily, Maidenhair Fern, Marigold, Narcissus, Palm, Pansy, Peony, Pelargonium, Petunia, Philodendron, Phlox, Pine, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulips, Viburnum, Wandering Jew, Yew, Zinnia and Andromeda, Arborvitae, Ash, Beech, Birch, Boxwood, Cotoneaster, Crabapple, Euonymus, Fir, Firethorn, Forsythia, Hawthorn, Hemlock, Hickory, Honey Locust, Horse Chestnut, Juniper, Larch, Laurel, Lilac, Linden, Mimosa, Myrtle, Oak, Pine, Privet, Tulip Tree, Viburnum, Willow.

Phytotoxicity:

Plant safety is an important consideration when using insecticides. However, it is not possible to evaluate the phytotoxicity of Evergreen® Pyrethrum Dust on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of Evergreen® Pyrethrum Dust treat a limited number of plants and observe for phytotoxicity over a 10 day period.

Evergreen® Pyrethrum Dust must come into contact with insects in order to control them. Coverage of the target insect is vital to achieving good performance. Apply Evergreen® Pyrethrum Dust to cover upper and lower surfaces of leaves stems and flowers with a thin, even film of dust. Apply when insects or insect damage first appears.

When treating trees, shrubs and similar plants, treat the trunk, the base of the trunk and the critical root zone. Treat the tree crotch, knotholes, wounds and other areas where insects may harbor. Allow dust to settle in treated areas before reentering. Do not water or irrigate immediately following treatment.

OUTDOOR AREAS for use in and around areas:

Ant Mounds	Decks	Gazebos	Outdoor Eating Facilities
Arboretums	Dog Runs	Home Gardens	Patios
Back Yards	Drive-In	Kennels	Porches
Balconies	Restaurants	Lanais	Sanitary Sewers (manholes)
Beneath Decks	Drive-In Theaters	Landscapes	Side Yards
Beneath Gazebos	Dumpsters	Lath Houses	Swine Yards
Calf Hutches	Farmers Markets	Loading Docks	Wood Piles
Compost Sites	Flea Markets	Lumber Storage	
Courtyards	Front Yards	Areas	
	Gardens	Nurseries	

Building Perimeter Treatments:

To aid in preventing entry and infestations of buildings by listed insects, Apply 16 to 24 oz. of Evergreen® Pyrethrum Dust per 1,000 square feet [4 to 6 ounces of Evergreen® Pyrethrum Dust per 250 square feet] to building exterior as a perimeter treatment in a band extending 12 - 18" along and adjacent to foundation, and to a height of 2 to 3 feet up the side of the building where these pests are active and may find entrance.

To Kill Insects Harboring in Lawns, Landscaping, and Backyards:

To aid in the control of listed insects, apply Evergreen® Pyrethrum Dust at the rate of 8 to 24 ounces per 1,000 square feet [2 to 6 ounces of Evergreen® Pyrethrum Dust per 250 square feet]. For best results, lawn should be mowed one to two (1 - 2) days before applying, and clippings removed. Be sure to cover the entire lawn surface where insects are present, or normally frequent. Apply under ornamental plants, shrubs and trees. Do not water or irrigate immediately following application. Keep pets and children out of the treated area until after the dust has settled.

LIVESTOCK, DAIRY, SWINE, POULTRY AND OTHER ANIMAL BUILDINGS AND PREMISES

WHEN USED ON DAIRY BARN OR FACILITIES: CLOSE MILK BULK TANK LIDS TO PREVENT CONTAMINATION FROM DUST AND FROM DEAD OR FALLING INSECTS. REMOVE OR COVER MILKING UTENSILS BEFORE APPLICATION. WASH TEATS OF ANIMALS BEFORE MILKING.

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Tick, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice, and Stable Flies

To aid in the control of listed insects in listed indoor and outdoor animal areas, apply Evergreen® Pyrethrum Dust at the rate of 1 oz. per 50 square feet to cracks, crevices, voids, structural joints, floors, walls, wall joints, support beams, trusses, rafters, ceilings, around scales, chutes, gates, and other areas where insects harbor, alight, hide or travel. For larger area treatments use 16 to 24 oz. of Evergreen® Pyrethrum Dust per 1,000 square feet [4 to 6 ounces of Evergreen® Pyrethrum Dust per 250 square feet]. Avoid contamination of feed, feedstuffs or milking equipment. For Ticks, apply Evergreen® Pyrethrum Dust, at the rate of 1 ounce per 70 square feet in places where the insects are traveling or harboring.

Apply 16 to 24 oz. of Evergreen® Pyrethrum Dust per 1,000 square feet [4 to 6 ounces of Evergreen® Pyrethrum Dust per 250 square feet] to exterior of livestock buildings in an 18" band along and adjacent to building foundations, and to a height of 2 to 3 feet up the side of the building where insects are active and may find entrance.

DOMESTIC ANIMAL USE**Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Tick, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice, and Stable Flies**

For Ticks, apply Evergreen® Pyrethrum Dust, at the rate of 1 ounce per 70 square feet in places where the insects are traveling or harboring.

Beef, Dairy Cattle, Horses and Other Livestock:

Can be used in dust bags, shaker can and mechanical dust applicator to aid in the control or repellency of listed livestock insects.

Direct Application: Apply 2 oz. of dust per animal by shaker can, or suitable mechanical dust applicator over the head, neck, shoulders, back and legs.

Application through Dust Bags: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

Swine:

Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Do not apply more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

Poultry:

To control listed livestock insects, apply at a rate of 1 lb. per 100 birds [10 ounces per 60 birds]. Ensure thorough treatment of vent area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Keep this product in its tightly closed original container, when not in use. Store in a cool, dry, well-ventilated (preferably locked) area that is inaccessible to children and animals, and away from food and pet food.

PESTICIDE DISPOSAL:

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry.)

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container. Completely empty container by shaking and tapping sides and bottom to loosen clinging particles promptly after emptying. Then offer for recycling if available or reconditioning, if appropriate or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with plenty of soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, Loaders Applicators, and other handlers must wear the following:

- long-sleeve shirt
- long pants
- shoes and socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

ENVIRONMENTAL HAZARDS

For terrestrial uses: This product is toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate. See Directions for Use for additional precautions and restrictions.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This pesticide is toxic to fish and other aquatic invertebrates. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

