

## FLEA, TICK & MANGE DIP

### FOR RESIDENTIAL USE ONLY

#### **ACTIVE INGREDIENT:**

Permethrin.....3.20% INERT INGREDIENTS: 96.80% TOTAL: ......100.00%

**KEEP OUT OF** CHII DREN

See Back Label Booklet for Additional Precautionary Statements

A CONTROL SOLUTIONS INC. PRODUCT

**NET CONTENTS** 





# FLEA, TICK & MANGE DIP FOR RESIDENTIAL USE ONLY

- Kills Fleas, Flies, Lice, Mange Mites and Ticks, including Deer Tick (carrier of Lyme Disease)
- Provides Contact and Residual Control for up to 4 weeks
- Makes 8 Gallons of Dip Solution

To Control: Flies, Fleas, Lice, Ticks on Dog and Dog Premises

Do Not Use On Puppies Less Than 12 Weeks of Age

#### ACTIVE INGREDIENT:

Permethrin	3.20%
INERT INGREDIENTS:	96.80%
TOTAL:	100.00%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Inside Booklet for Additional Precautionary Statements

Control Solutions, Inc. 5903 Genoa-Red Bluff • Pasadena, TX 77507-1041 281-892-2500

# READ ENTIRE LABEL BEFORE USE FOR PET USE - USE ONLY ON DOGS

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

#### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall<sup>®</sup> International (866) 897-8050 for emergency medical treatment information.

#### If swallowed

- Call poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
    Do not induce vomiting unless told
- to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN AND VETERINARIANS:** Do not administer or instill milk, cream or other substances containing vegetable or animal fats which will enhance absorption of lipophilic substances.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively, visiting the treatment area.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use in undiluted form.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep people and pets off treated areas until the spray has dried.

#### SHAKE WELL BEFORE USING

Use Dilution rate as indicated: 1 fl. oz. = 6 tsp. or 2 Tbsp. 8 fl. ozs. = 1/2 pt. or 1 cup

#### DOG AND DOG PREMISE USE - REFERENCE CHART

DILUTIONS PESTS (Mix Well)

HOW TO APPLY

## PESTS ON DOG PREMISES (Dog Houses, Runs and Yards)

Fleas, Flies, Dilute 4 fl. Thoroughly spray infested areas, pet beds and resting quarters. Fresh bed-Ticks, Lice and ozs. in 1 gal. ding should be placed in animals quarters following treatment. Do not spray Mange Mites feed, food or water. Repeat as needed but not more than once every two water (adult and larvae) weeks.

Fleas, Ticks, Dilute 2 fl including Deer ozs, in 1 gal. tick (carrier of water Lyme disease). Lice & Mange

## PESTS ON DOGS (DO NOT TREAT PUPPIES LESS THAN TWELVE [12] WEEKS OF AGE) Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to

two to three weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

dry in a warm place without rinsing or toweling. Do not reapply product for

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. Store in a locked area out of reach of children and domestic animals. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: If empty - Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

If partly filled: Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

**NOTE:** Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.





## FLEA, TICK & MANGE DIP

- READ ENTIRE LABEL BEFORE USE
- FOR PET USE USE ONLY ON DOGS

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

#### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

## If swallowed

- Call poison control center or doctor immediately for treatment advice.
   Have person sip a glass of water if able to swallow
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN AND VETERINARIANS:** Do not administer or instill milk, cream or other substances containing vegetable or animal fats which will enhance absorption of lipophilic substances.

#### CONTROL SOLUTIONS, INC.

5903 Genoa-Red Bluff • Pasadena, TX 77507-1041 281-892-2500

EPA Reg. No. 53883-78 EPA Est. No. 53883-TX-002

0411