

# TOP GUN™

# SPECIMEN LABEL

## All-Weather BAIT BLOCK® Rodenticide Contains BITREX® for Indoor and Outdoor Rodent Control

**Kills Norway Rats,  
Roof Rats & House Mice.**

House Mice, Norway Rats and Roof Rats usually consume a lethal dose in a single night's feeding, but it may take two or more days from time of bait consumption for these rodents to die. Rodents cease feeding after consuming a toxic dose. This product is effective against anticoagulant-resistant mice and rats.

**ACTIVE INGREDIENT:**

Bromethalin CAS [63333-35-7] ... 0.01%

**OTHER INGREDIENTS:** ..... 99.99%

**TOTAL:** 100.00%

EPA REG. NO. 67517-66-56

EPA EST. NO. 56-OH-1

1/2 OZ. BAIT BLOCK®

### KEEP OUT OF REACH OF CHILDREN CAUTION

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**MAY BE HARMFUL IF EATEN BY CHILDREN OR PETS**

Manufactured By: J.T. Eaton & Co., Inc.,  
1393 East Highland Road, Twinsburg, OH 44087 U.S.A.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Keep away from humans, domestic animals and pets. May be harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling bait.

#### FIRST AID

##### HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

**IF SWALLOWED** • Call a Poison Control Center, doctor or \*HOT LINE NUMBER 1-800-424-9300 immediately for treatment advice.  
• Have person sip a glass of water if able to swallow.  
• Do not induce vomiting unless told to do so by the poison control center or doctor.

**IF ON SKIN OR CLOTHING** • Take off contaminated clothing.  
• Rinse skin immediately with plenty of water for 15-20 minutes.  
• Call a poison control center, doctor or \*HOT LINE NUMBER 1-800-424-9300 immediately for treatment advice.

You may also contact **HOT LINE NUMBER 1-800-858-7378 National Pesticide Information Center (NPIC)** for health concerns, emergency medical treatment information or pesticide incidents.

##### NOTE TO PHYSICIAN:

This product is not an anticoagulant type of rodenticide. If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

**IN CASE OF EMERGENCY CALL : CHEMTREC (POISON CONTROL CENTER) at 1-800-424-9300**

### ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

### STORAGE AND DISPOSAL

#### [Household Use:]

**STORAGE:** Store only in original container, in a cool, dry place inaccessible to children and pets.

**DISPOSAL: If Empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or **1-800-CLEANUP** for disposal instructions. Never place any unused product down any indoor or outdoor drain.

#### [Non-household Use:]

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store only in original container, in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** This product may be used to control Norway rats, roof rats, and house mice in sewers and for use inside and against outside walls of homes, industrial and agricultural buildings and similar man-made structures. This product may also be used in alleys located in urban areas, inside transport vehicles (ships, trains, aircraft), and inside and against outside walls of related port or terminal buildings. For use in non-food/non-feed areas. Do not contaminate human or pet food preparations items or areas. Do not place near or inside ventilation duct openings. Do not broadcast bait.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where rodents will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Remove as much alternative food accessible to rodents as possible.

#### APPLICATION DIRECTIONS:

The active ingredient in this product is different from anticoagulant rodenticides. Individual rodents cease feeding after consuming a toxic dose.

**Rats:** Apply 2 to 12 baits per placement. Space placements at intervals of 20 to 30 feet.

**Mice:** Apply 1 or 2 baits per placement. Space placements at intervals of 8 to 12 feet.

**FOLLOW-UP:** Adjust the amount of bait applied to the level of rodent feeding expected at each bait placement location. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect bait placements and replenish bait at sites where there is heavy feeding or where there is continued evidence of rodent activity. Maintain bait for at least one week and continue baiting until all signs of rodent activity have ceased or it becomes clear that the remaining rodents are not attracted to the bait. Collect and properly dispose of all dead rodents and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

#### Notice of Warranty

J.T. EATON & CO., INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

Manufactured by: J.T. EATON & CO., INC., 1393 East Highland Road  
Twinsburg, OH 44087 U.S.A. ©2005

BAIT BLOCK® is a registered trademark and JT EATON™ and TOP GUN™ are trademarks of the J.T. Eaton & Co., Inc., Twinsburg, OH 44087  
BITREX® is a registered trademark of Macfarlan Smith, Scotland  
10.20.2003/02.04.2005