

EIGHT® INSECT CONTROL YARD & GARDEN READY TO SPRAY

CONTROLS: ANTS, APHIDS, ASIAN LADYBUGS, BAGWORMS, BEETLES, BORERS, TICKS, CHINCH BUGS, CLUSTER FLIES, CODLING MOTHS, CRICKETS, EARWIGS, FLEAS, CICADAS (LOCUST), GNATS, GRASSHOPPERS, JAPANESE BEETLES, SCALES, THRIPS, WEEVILS, WHITEFLIES, MOLE CRICKETS, SPIDER MITES, TENT CATERPILLARS, MEALYBUGS, LEAFMINERS, AND MANY OTHER LISTED INSECTS.

FOR USE ON LISTED: LAWNS, TREES, SHRUBS, FLOWERS, FRUIT, NUTS, VEGETABLES, OUTSIDE SURFACES OF BUILDINGS AND SURROUNDING AREAS.

APPLY USING THIS HOSE END APPLICATOR.

NO MIXING REQUIRED.

EPA Est. No. 4-NY-1

EPA Reg. No. 4-407

Buyers Guarantee Limited to Label Claims.

Bonide Products, Inc.

6301 Sutliff Road

Oriskany, NY 13424

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

For Residential use only.

SHAKE WELL BEFORE USING

HOSE END SPRAYER: EASY TO USE

Make sure knob Control lever is in the OFF position. Shake well and attach to hose. Turn water on at faucet, aim nozzle toward lawn and turn knob Control lever to ON to begin spraying. When finished turn Control lever to OFF position. Turn faucet to OFF position. Discharge residual water pressure by turning ON and OFF again. Store out of reach of children.

People and pets should not be allowed in treated areas until spray has dried.

DIRECTIONS FOR APPLYING HOME LAWN

| | Pests: | How to Use: |
|--|--|--|
| Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine | Ants, Armyworms, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Fleas, Grasshoppers, Japanese Beetle Grubs, Mole Crickets, Sod Webworms, Ticks that may carry Lyme Disease, Leafhoppers | Home lawns should be 3 inches or less in height at the time of application. 1 Quart of this concentrate as applied through the Hose-End Sprayer treats 5,000 sq. ft. For heavy infestations, repeat application after 2 weeks. |

TREES & SHRUBS

| | Pests: | How to Use: |
|--|---|--|
| Arizona Cypress, Azalea, Birch, Cherry, Non-Bearing Citrus, Conifers, Elm, English Ivy, Euonymous, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-Orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar, Rhododendron, Taxus | Aphids, Bagworms, Cicadas (Locust), Exposed Thrips, Fall Cankerworms, Inchworms, Leafminers, Leafrollers, Locust, Mealybugs, Spider Mites, Whiteflies, Gypsy Moth, Tent Caterpillar, Japanese Beetles | Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use at intervals of 4-8 days. Application can be made up to the day of collection. |
| Cherry, Honeysuckle | Fall Webworms | |

TREES & SHRUBS Cont.

| | Pests: | How to Use: |
|------------------------|--|--|
| Conifer | Budworms | Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° F to 75° F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use at intervals of 4-8 days. Application can be made up to the day of collection. |
| Elm | Elm Leaf Beetles, Elm Spanworms, Fall Webworms | |
| Douglas and Grand Firs | Tussock Moths | |
| Non-bearing Citrus | Citrus Black Flies | |
| Oak | Oakworms | |
| Pine | Pine Beetles, Pine Moths, Pine Needleminers, Needle Scales | |

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

ROSES & FLOWERS

| | Pests: | How to Use: |
|--|--|---|
| Ageratum, Ardisia, Aster, Azalea, Baby's Breath, Begonia, Birds Nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia | Aphids, Armyworms, Bagworms, Cabbage Worms, Cicadas (Locust), Cornworms, Cutworms, Exposed Thrips, Fall Cankerworms, Fall Webworms, Fungus Gnat, Inchworms, Japanese Beetles, Leafminers, Leafrollers, Locust, Mealybugs, Scale Crawlers, Spring Cankerworms, Spider Mites, Whiteflies, Rose Chafer, Psyllids, Scale Insects | Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° F to 75° F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection. |

FRUIT & NUT TREES

GENERAL DIRECTIONS- Apply when insects appear. Wet plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

| | Pests: | How to Use: |
|--|--|---|
| Almonds, Pecans | Navel Orange Worms, Peach Twig Borers. | Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest. |
| Apples | Green Fruitworms, Aphids, Leafrollers (Oblique-banded, Redbanded), Plum Curculio, Rosy Apple Aphids, Spotted Tentiform Leafminers, Tarnished Plant Bugs, White Apple Leafhoppers, Japanese Beetles | Repeat as required to maintain control. Do not make more than 3 applications. Do not apply after petal fall. |
| Peaches | Borers (Lesser Peachtree, Peach Twig), Green Fruitworms, Plum Curculio, Oriental Fruit Moths, Tarnished Plant Bugs. | Do not apply within 7 days of harvest. Do not make more than 8 applications per season. |
| Pears (dormant through delayed dormant) | Pear Psylla | Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season. |
| Pears (summer) | Codling Moths, Green Fruitworms, Pear Psylla, Aphids | Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest. |

VEGETABLES

| | Pests: | How to Use: |
|-----------------------------|---|--|
| Asparagus (pre-harvest) | Asparagus Beetles, Cutworms | Apply as needed to ensure uniform coverage. Do not make more than 4 applications per crop. Can be applied up to 3 days prior to harvest. |
| Asparagus (post-harvest) | Larval stages of Asparagus Beetle; Tarnished Plant Bug, Lygus Bugs and Adult Japanese Beetle. | Treat ferns or bush growth after spear harvest when insects are present. |

VEGETABLES Cont.

| | Pests: | How to Use: |
|--|---|---|
| Celery—see Leafy Vegetables | | |
| Cole Crops, Broccoli, Brussels Sprouts, Cabbage, Cauliflower | Beet Armyworms, Cabbage Aphids and Loopers, Diamondback Moths, Imported Cabbage Worms | Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli and 10 applications on Cabbage. |
| Corn – see Sweet Corn | | |
| Cucurbits: Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon | Aphids, Cabbage Looper, Cucumber Beetle (adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant Bugs, Squash Bugs. | Apply as needed to ensure uniform coverage. Can be applied up to the day of harvest. |
| Eggplant | Colorado Potato Beetles, Vegetable Leafminers | Apply as needed to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest. |
| Horseradish (Foliar application) | Imported Crucifer Weevils | Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest. |

VEGETABLES Cont.

| | Pests: | How to Use: |
|---|--|---|
| Leafy Vegetables: Amaranth, Arugula, Celery, Celtuce, Chervil, Corn salad, Chrysanthemum (edible-leaved and garland), cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Okra, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard | Alfalfa Looper, Aphids, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworm, Southern Armyworm, Tobacco Budworm, Vegetable Leaf Miner. | Apply every 5 to 10 days as needed. Do not make more than 7 applications per crop. Can be applied up to 1 day prior to harvest. |
| Melons – see Cucurbits | | |
| Potatoes | Beet Armyworm, Colorado Potato Beetle, Potato Leafhopper, Potato Tuberworm, Cabbage Looper, Cutworms, European Corn Borer, Potato Psyllid, Tarnished Plant Bug. | Apply as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest. |
| Peppers | Vegetable Leafminers, Cabbage Looper, Flea Beetle, Corn Earworm, Pepper Weevil, European Corn Borer. | Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest. |
| Strawberries, Raspberries, Blueberries | Strawberry Weevils, Aphids, Spider Mites, Lygus Bugs, Flea Beetles, Omnivorous Leafhopper, Spittlebugs, Japanese Beetles, Field Crickets, Red Spider, Leafroller, Weevils. | Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season. |
| Sweet Corn | Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm. | Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest. |
| Tomatoes | Beet and Southern Armyworm, Tomato Pinworm, Tomato Fruitworm, Horn Worms, Granulate Cutworms, Colorado Potato Beetle, Cabbage Looper | Apply as needed for uniform coverage. Can be applied up to day of harvest. |

NUISANCE PESTS IN OUTDOOR AREAS

GENERAL DIRECTIONS

Activate Hose-End Sprayer and spray mist listed areas for thorough coverage.

OUTSIDE SURFACES OF BUILDING

Use this product on Ants, Clover Mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

OTHER OUTDOOR AREAS

Use this product on Ants, Asian Ladybugs, Crickets, Earwigs, Fleas, Locust, Ticks (**including ticks that may carry Lyme Disease**) for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For Ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

To reduce annoyance from: **Cluster Flies, Flying Moths, Gnats, House Flies, Mosquitoes.** Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate. Repeat as necessary.

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Disposal: *If empty* - Do not reuse this container. Place in trash or offer for recycling if available.
If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

| | |
|--------------------------------|---|
| If Swallowed: | <ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If on Skin or Clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| If Inhaled: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |
| If in Eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. Avoid contamination of ornamental fish ponds. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.