



ferti·lome®



# WEED FREE ZONE

READY-TO-SPRAY

*Clover*

*Ground Ivy*

*Spurge*

- Rainfast (Rainproof) in 3 hours.
- Proven Cool Weather Performance.

**THIS PRODUCT CONTAINS:**

0.52 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.96%.  
0.20 lb. (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 2.66%.

0.05 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.67%.

0.04 lb. Ethyl  $\alpha$ ,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate or 0.54%

Isomer Specified by AOAC Methods.

Contains Petroleum Distillate

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**ACTIVE INGREDIENTS:**

2,4-D, 2-ethylhexyl ester ..... 10.49%

Mecoprop-p, acid ..... 2.66%

Dicamba, acid ..... 0.67%

Carfentrazone-ethyl ..... 0.54%

**INERT INGREDIENTS:** ..... 85.64%

**TOTAL** ..... 100.00%

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

See Back Panel & Booklet  
for First Aid, Precautionary  
Statements and  
Directions for Use.

**NET CONTENTS ONE QUART (32 FL. OZS.)**

ferti-lome® Weed Free Zone RTS

Produced For:  
VOLUNTARY PURCHASING GROUPS, INC.  
230 FM 87  
BONHAM, TEXAS 75418  
Visit Us At: [www.fertilome.com](http://www.fertilome.com)  
EPA Reg. No. 2217-855-7401  
EPA Est. No. 7401-TX-1



**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

***Personal Hygiene Statements:***

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**FIRST AID**

<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a Poison Control Center or doctor immediately for treatment advice.</li><li>• Do not induce vomiting unless told to by a Poison Control Center or doctor.</li><li>• Do not give any liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If On Skin:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a Poison Control Center or doctor for treatment advice.</li></ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

You May Also Contact 1-877-800-5556 for emergency medical treatment advice.

**NOTE TO PHYSICIAN:** Contains petroleum distillate—vomiting may cause aspiration pneumonia.

PEEL FROM CORNER OF BOOKLET  
OPEN RESEALABLE LABEL FOR DIRECTIONS & PRECAUTIONS • PRESS TO RESEAL

5W-4-49H

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsates.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

## PRODUCT DESCRIPTION

This Trimec® product is a combination of four proven Weed Killers: 2,4-D, mecoprop-p, dicamba and carfentrazone-ethyl. Postemergent applications of this product control a wide range of Broadleaf Lawn Weeds in Residential Lawns including Dandelion, Chickweed, Spurge, Pennywort (Dollarweed), Knotweed, Plantain, Clover and Ground Ivy (Creeping Charlie).

This product offers these advantages:

- Rapid and effective Weed control for common and troublesome Weed species in turf grass.

- Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed within hours of the application and plant death can occur within 7 - 14 days.

#### WHERE TO USE

This product is intended for use on most Ornamental Lawns including:

COOL SEASON TURFGRASS	WARM SEASON TURFGRASS
Kentucky Bluegrass	Common Bermudagrass
Perennial Ryegrass	Hybrid Bermudagrass
Tall Fescue	Bahia grass
Red or Fine Leaf Fescues	Zoysiagrass
Colonial Bentgrass	Buffalograss
Mixtures of Cool Season Species	St. Augustinegrass
	Centipedegrass

#### **Turfgrass Tolerance:**

The turfgrass tolerance to **ferti-lome® Weed Free Zone** may vary and temporary turfgrass yellowing may occur on St. Augustinegrass and certain varieties of Hybrid Bermudagrass.

**ferti-lome® Weed Free Zone** may injure certain turfgrass species. To reduce risk of Lawn injury, avoid spraying drought stressed Lawns or when temperatures are above 90°F.

#### **Prohibitions:**

- Do not use this product on 'Floritam' or 'Bitterblue' varieties of St. Augustinegrass in Florida.
- Do not use this product on Carpetgrass, Dichondra or Lawns containing desirable Clovers.
- Do not apply this product to St. Augustinegrass during Spring green-up which is the transition period between dormancy and active growth.

- Do not spray roots of Trees or Ornamentals.
- Do not allow spray drift onto Vegetables, Flowers, Ornamental Plants, Shrubs, Trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Be particularly careful within drip-line of Trees and other Ornamental species.
- Do not apply as a fine mist because of potential injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product runoff, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

#### WHEN TO APPLY

Spray when Broadleaf Weeds are young and actively growing for best results.

**ferti-lome® Weed Free Zone** may be applied in the Spring, Summer, or Fall. Spring and Fall treatments under adequate soil moisture conditions are preferred to the Summer treatments. Generally, Summer broadcast application to older, drought stressed Weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2-6 week intervals are recommended for more mature Weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations or as a follow-up treatment, or any time broadleaf weeds are susceptible.

#### CULTURAL TIPS FOR IMPROVED CONTROL

##### ***Irrigation:***

- Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

##### ***Mowing:***

- Delay mowing 1 to 2 days before and after the application of this product.

##### ***Reseeding Interval:***

- Treated areas may be reseeded 2 weeks after application.

**RATE RECOMMENDATIONS FOR COOL SEASON TURFGRASS.**

<b>Species</b>	<b>Amount of Product</b>	<b>Treated Area, sq. ft.</b>
	<b>fl. oz./1,000 sq. ft.</b>	<b>1.0 qt. (32 fl. oz.) Sprayer Jar or Retail Container</b>
Kentucky Bluegrass	1.5 to 2.0	16,000 to 21,300 sq.ft
Perennial Ryegrass		
Tall Fescue		
Red or Fine Leaf Fescues		
Colonial Bentgrass		
Mixtures of Cool Season Species		

**RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS.**

<b>Species</b>	<b>Amount of Product</b>	<b>Treated Area, sq. ft.</b>
	<b>fl. oz./1,000 sq. ft.</b>	<b>1.0 qt. (32 fl. oz.) Sprayer Jar or Retail Container</b>
Common Bermudagrass	1.0 to 2.0	16,000 to 32,000 sq. ft.
Hybrid Bermudagrass	0.75 to 1.5	21,300 to 43,000 sq.ft.
Bahiagrass		
Zoysiagrass		
Buffalograss		
St. Augustinegrass		
Centipedegrass		

<sup>1</sup>Do not apply this product to 'Floritam' or 'Bitterblue' varieties of St. Augustinegrass in Florida.

The maximum application rate is 3.0 fl.oz. of product/1,000 sq.ft. per application (0.5 lbs 2,4-D acid equivalent per acre per application). The maximum number of

broadcast applications is limited to 2 per year. The maximum seasonal rate is 6.0 fl.oz. of product/1,000 sq.ft. (1.0 lb 2,4-D acid equivalent per acre), excluding spot treatments.

### **How To Apply Using The Ready To Spray System**

1. Measure the total (entire) lawn to be sprayed. For rectangle areas, multiply the length by the width to obtain total square feet.
2. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the OFF position.

TO BEGIN spraying: Turn on water at faucet. Point nozzle toward lawn to be treated and turn the control valve to the ON position. water will automatically mix with the product. Spray evenly over the measured area.

3. SPRAY: spray evenly over the measured area. Walk at a steady pace. Refer to the view strip [graduated scale] on the side of the container to guide coverage. The 32 fl.oz. container will treat 16,000 to 43,000 sq.ft. of turfgrass.
4. To STOP spraying: Turn the control valve to the OFF position. turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

### **BROADLEAF WEEDS CONTROLLED**

**ferti-lome® Weed Free Zone** will control or suppress the following Broadleaf Weeds and will control or suppress other Broadleaf Weeds that are susceptible to 2,4-D. A partial list is presented on the following page.

### BROADLEAF WEEDS

Aster, White Heath And White Prairie	(*Creeping Oxeye)	Poison Ivy
Bedstraw	Filaree, Whitestem And Redstem	Poison Oak
Beggarweed, Creeping	Florida Betony	Prostrate Knotweed (*Knotweed)
Bindweed	Florida Pusley	Puncturevine
Black Medic	Ground Ivy (Creeping Charlie)	Purple Cudweed
Broadleaf Plantain	Groundsel	Purslane
Buckhorn Plantain	Hawkweed	Ragweed
Bull Thistle	Healall	Redweed
Burclover	Henbit	Red Sorrel (*Sheep Sorrel)
Burdock, Common	Innocence	Shepherdspurse
Buttercup, Creeping	(Blue-eyed Mary)	Spotted Spurge
Carpetweed	Lambsquarters	Spurge
Chickweed, Common	Lawn Burweed	Thistle
Chicory	Lespedeza, Common	Veronica
Cinquefoil	Mallow, Common	(*Corn Speedwell)
Clover	Matchweed	Virginia Buttonweed
Compassplant	Mouseear Chickweed	White Clover
Curly Dock	Old World Diamond	(*Dutch Clover,
Dandelion	Flower	Honeysuckle Clover,
Dayflower	Oxalis (*Yellow Wood- sorrel And	White Trefoil And Purplewort)
Deadnettle	Creeping Woodsorrel)	Wild Carrot
Dock	Parsley-piert	Wild Garlic
Dogfennel	Pennsylvania	Wild Geranium
English Daisy	Smartweed	Wild Lettuce
False Dandelion	Pennywort	Wild Mustard
(*Spotted Catsear And Common Catsear)	(*Dollarweed)	Wild Onion
Field Bindweed	Pepperweed	Wild Strawberry
(*Morningglory And Creeping Jenny)	Pigweed	Wild Violet
Field Oxeye-daisy	Pineappleweed	Yarrow
	Plantain	Yellow Rocket

\*Synonyms



## STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**CONTAINER DISPOSAL:** **If Empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If Partially filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the Seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

